


AMRO/PAHO Region

Fentanyl Consumption (mg/person), 1985 - 2020



AMRO/PAHO Fentanyl Consumption, (mg/person)

| Year | Fentanyl |
|------|----------|
| 1985 | 0.0067 |
| 1986 | 0.0076 |
| 1987 | 0.0074 |
| 1988 | 0.0114 |
| 1989 | 0.0079 |
| 1990 | 0.0149 |
| 1991 | 0.0534 |
| 1992 | 0.0585 |
| 1993 | 0.0573 |
| 1994 | 0.0846 |
| 1995 | 0.1011 |
| 1996 | 0.1020 |
| 1997 | 0.1165 |
| 1998 | 0.1444 |
| 1999 | 0.1717 |
| 2000 | 0.2183 |
| 2001 | 0.2538 |
| 2002 | 0.3389 |
| 2003 | 0.4535 |
| 2004 | 0.5306 |
| 2005 | 0.6445 |
| 2006 | 0.7626 |
| 2007 | 0.7752 |
| 2008 | 0.8578 |
| 2009 | 0.7369 |
| 2010 | 0.6481 |
| 2011 | 0.7320 |
| 2012 | 0.6243 |
| 2013 | 0.6551 |
| 2014 | 0.5953 |
| 2015 | 0.5955 |
| 2016 | 0.5501 |
| 2017 | 0.3933 |
| 2018 | 0.3800 |
| 2019 | 0.2138 |
| 2020 | 0.2672 |

Data sources:

Consumption Data – International Narcotics Control Board

Population – The World Bank, United Nations Statistics Division

ME Conversion Factors – WHOCC Centre for Drug Statistics Methodology

Walther Global Palliative Care
& Supportive Oncology
Indiana University
Melvin and Bren Simon
Comprehensive Cancer Center